



ProFouND: Prevention of Falls Network for Dissemination

DELIVERABLE D 4.4

Updated EVIDENCE SYNTHESIS AND GENERIC GUIDANCE

Document Type:	Deliverable
Dissemination Level:	check status
Editor:	RBMF
Document state:	Final
Document version:	1.1
Contributing Partners:	UNIMAN, GCU, LLT
Contributing WPs:	WP4, WP1 WP3, WP5, WP8
Estimated P/M (if applicable):	1
Date of Completion:	21/10/2015
Date of Delivery to EC:	21/10/2015)
Number of Pages:	40

Table of content

1	Description of deliverable, tasks and Milestones for WP4	3
2	Development of deliverable	4
3	Summary and Perspectives	5
4	Appendix Updated Factsheets	6

1 Description of deliverable, tasks and Milestones for WP4

Objectives:

A central objective of ProFouND has been to construct a system for providing customised information to users, be they health care provider organisations, health or social care professionals, NGOs, or older people themselves.

WP4 has provided a first version of the evidence and generic guidance in month 19. This will be done by constructing a library of evidence for each domain based on reviews of evidence. These factsheets have been disseminated via direct action to different stakeholders, via the website and extensive support from members and associates of the consortium. Comments, advice and criticism has been received and was discussed within the consortium. The literature has been continuously reviewed to include new studies and review papers.

The deliverable for WP4 month 31 was the following:

D4.4 Updated evidence synthesis and generic guidance [Month 31]

WP4 has performed the following task:

Task 4.1: WP4 has used existing systematic reviews and evidence syntheses to identify and operationalize best practice in requisite areas and put specific attention onto how to implement evidence and overcome barriers to innovation in practice.

We have started in areas where the evidence is strong and relatively clear cut, regarding strength and balance exercises, medication review, environmental modification for specific high risk groups. In these cases we have presented specific recommendations of best practice and guidance of how to implement this in practice.

Where evidence is poorer we were possible to undertake our own review and synthesis.

We knew from reviews already undertaken that the evidence base for ICT and other technologies being effective in falls prevention/detection is emerging is still limited.

This updated guidance is aiming at patient interest groups, policy makers, health care funders as well as other professional groups [M19-M31].

WP4 has reached the following Milestones between months 19 and 31:

MS24 network consensus meeting on evidence synthesis

2 Development of deliverable and updated evidence synthesis and generic guidance

During the past 12 month the evidence and guidance has been successfully disseminated, distributed and more importantly used in several regions and countries. This processed is based on the successful co-operation with partners, associates, with groups of the EIP AHA 2 initiative, stakeholder organizations such as the SIG of EUGMS, NGOs and governmental organization.

Examples from Germany include the Bundeszentrale für Gesundheitliche Aufklärung (BZgA) as a branch of the German Ministry of Health (BMG) which started a campaign on fall prevention based on ProFouND material after a visit of the WP leader and PI Prof. Todd in May 2014. In July 2015 this led to web based material, printed brochures and poster material for GP officies and pharmacies across Germany. More than 20,000 copies of the brochures were distributed in the first 3 month after the release (<http://www.gesund-aktiver-werden.de/>). The health care fund for farmers that is crucial for rural areas has also started a campaign on fall prevention in rural areas in Germany for several 10,000 older persons. In this campaign the ProFouND material in its German version is fully included (www.trittsicher.org). Many other examples can mentioned were the guidance and the summarized evidence have been stimuli and facilitators for dissemination campaigns.

As expected we have received not only positive feedback but also recommendations for improvement. A lot of these comments were based on personal experience and observation, sometimes on local structures leading to barriers to implementation. Sometimes very committed experts also reported frustration on the slow speed of social innovations. This encouraged us to include these comments as much as possible for the final dissemination event of ProFouND in Bologna in Feb 2015.

Valuable comments were received from our advisory board and non-European international experts. Most of these led to minor changes of the fact sheets now documented in a printable format.

3 Summary and perspectives

During the remaining time of the network WP 4 is aiming to reach many stakeholder organisations, policy makers and health care managers to inform them about best evidence, implementation needs and current developments. This will be done in close co-operation with the other WPs. A final highlight of this effort will be the next European Falls Festival where relevant persons and organizations will be reached.

The updated Cochrane reviews in the domains of community dwelling older adults, long-term and acute care as well as community approaches were expected were expected for late 2015 but might be delayed to be included in the final document in January 2016. We will be contact with the working groups of the Cochrane consortium.

The work on fact sheets on fall prevention in long-term care and acute care has started. This is stimulated by recent publications in the Lancet reporting on innovative successful approaches in settings that were thought to be difficult to be included in this ubiquitous problematic area.

A final aim for the upcoming update is the description of research areas in this domain where observational studies indicate room for improvement but randomized trials are still lacking. Examples are material science approaches for floor material or improved designs of walking aids and wheelchair ergonomics.



Falls Prevention Intervention Factsheets



Content

1.	General guidance	2
<hr/>		
2.	Exercise	9
<hr/>		
3.	Vision	15
<hr/>		
4.	Bone health	18
<hr/>		
5.	Vitamin D	23
<hr/>		
6.	Home & environment	26
<hr/>		
7.	Footwear & protective clothing	30
<hr/>		
8.	Falls detection & prevention technologies	33
<hr/>		



Generic guidance and implementation guidelines

Implementation Guidance

Target group:

Older people living in the community (see individual factsheets alongside this guidance).

Multifactorial interventions:

Multifactorial interventions assess an individual’s risk of falling, and then carry out individualised treatment or arrange referrals to reduce the identified risks. Multifactorial interventions often comprise the components described below. They are complex interventions and need to be carried out by specific health care professionals working together in a multidisciplinary process. Overall, current evidence shows that this type of intervention reduces the number of falls but not the number of people falling. This suggests that they are particularly recommended for people who have recurrent falls.

Exercise:

Selected group and home-based exercise containing progressive balance exercises and strength training effectively reduce the number of falls and the number of people falling. The programmes with the best evidence are the Otago Exercise Programme (OEP), Tai Chi, and the Falls Management Exercise programme (FaME -sometimes called PSI). Evidence suggests that new programmes such as Lifestyle integrated Functional Exercise (LiFE) might achieve similar or even better results.

Exercise alone has been demonstrated to be effective for persons after a first fall /with lower risk of falling. Overall, exercise interventions reduce fall-related fractures. Exercise must be challenging, progressive and regular, last more than 50 hours, and the regimen should be adhered to in the long term to be effective.

Medication and medical intervention:

Taking vitamin D supplements may be effective in reducing falls in people with low vitamin D levels in the blood before treatment. This needs to be carefully discussed with the responsible physicians.

Some medications increase the risk of falling. Gradual withdrawal of psycho-tropic medication (e.g. benzodiazepines, Zolpidem) for improving sleep, reducing anxiety and treating depression has been shown to reduce falls.

Insertion of a pacemaker can reduce falls in people with frequent falls associated with carotid sinus hypersensitivity and cardiac dys-/arrhythmias. Treatment of specific underlying causal factors, specific therapy, medication reviews and prescription modification programmes can also be effective in reducing the risk of falling, e.g. in people who fall because of hypervolemia (restoring haemodynamic stability) or orthostatic hypotension (discontinuing medication).

Home and environment:

Interventions to improve home safety appear to be effective for people at high risk of falling, particularly those with severe visual impairment. They should be carried out by suitable qualified health care professionals- usually occupational therapists.

Feet and Footwear:

Feet and Footwear assessments followed by podiatric care, foot and ankle exercises and provision of appropriate footwear can reduce the number of falls in people with disabling foot pain.

Vision and vision aids:

Adjustment of optical aid / spectacles (wearing single lens instead of multifocal glasses when outdoors if already active outdoors) and cataract surgery can reduce falls.

Implementation guidelines for fall prevention interventions:

Falls are everyone's business as there are multiple factors associated with them. Therefore establishing links between acute, community and primary health and social care services, the voluntary sector, charities and private companies is very important.

It is beneficial to assess the older person in their own environment as you are more likely to identify underlying problems and understand their needs.

If the team assessing the older person also carries out the interventions, success and uptake is more likely.

Follow-up any referrals to other services to ensure action has been taken, communication with all other services is important.

Further information (e.g. leaflets) should be discussed with the older person and personalised to their needs (not just handed out).

One size does not fit all. Ensure that the older person knows that the intervention is tailored to their needs. They are more likely to be motivated to comply with it.

Older people may have specific goals they want to achieve. Assisting them to set these goals and then work towards them will help them understand the importance of the changes they want to make and increase their confidence.

Older people can be easily put off engaging in a falls prevention programme if something goes wrong e.g. transport does not turn up, they were not told a session was cancelled. Ensure good communication is maintained at all times.

Engage family members and friends where possible as these can be a key source of support.

Give feedback to the older person's general practitioner, as positive affirmation from them can support action from the patient.

Ensure you follow the older person up either by phone call or preferably in person to monitor their progress. Give feedback on their progress, highlighting what they have achieved, it will help with motivation and it will also provide evidence for commissioners (funders of services).

When promoting your service focus on the positive action that older people can take to promote healthy ageing, rather than falls prevention itself.

Here are some suggestions:

If an older person is highly fearful of falls and pro-actively seeks advice about preventing falls then you can talk to them about risk management and prevention of falls. For all other older people do not initially talk about risk and reducing their risk of falling, they are unlikely to identify themselves as being at risk. You may want to invite them to an 'ageing well' assessment rather than a 'falls' assessment.

Instead talk about:

- How strength and balance training will improve their function e.g. Getting up and down stairs more easily and how it will help them to maintain their independence.
- How practising techniques for getting down and up from the floor means that they can play with their grandchildren.
- How being given a full health assessment and tailored advice gives them the knowledge and opportunity to take control of the situation and promote their own health and well-being (for the future if they do not see an issue now).

- How interventions you can offer will help them to manage their health conditions e.g. osteoporosis, arthritis, Parkinson's, stroke.
- Consider getting other older adults who have had positive experiences of your service to share their thoughts and support others, peer support can be very effective.

What did not work?

There is no evidence of an effect of cognitive behavioral interventions on the rate of falls. Trials testing interventions to increase knowledge and educate about fall prevention alone did not significantly reduce the rate of falls.

Links and Resources

profound.eu.com

References

Dorgo, S., King, G. A., Bader, J. O., & Limon, J. S. (2011). Comparing the effectiveness of peer mentoring and student mentoring in a 35-week fitness programme for older adults. *Archives of Gerontology and Geriatrics*, 52(3), 344–349. doi:10.1016/j.archger.2010.04.007

Gates, S., Fisher, J. D., Cooke, M. W., Carter, Y. H., & Lamb, S. E. (2008). Multifactorial assessment and targeted intervention for preventing falls and injuries among older people in community and emergency care settings: systematic review and meta-analysis. *British Medical Journal (Clinical Research Ed.)*, 336(7636), 130–133. doi:10.1136/bmj.39412.525243.BE

Gillespie, L. D., Robertson, M. C., Gillespie, W. J., Sherrington, C., Gates, S., Clemson, L. M., & Lamb, S. E. (2012). Interventions for preventing falls in older people living in the community. *The Cochrane Database of Systematic Reviews*, 9, CD007146. doi:10.1002/14651858.CD007146.pub3

Haran, M.J., Cameron, I.D., Ivers, R.Q., Simpson, J.M., Lee, B.B., Tanzer, M., Porwal, M., Kwan, M.M.S., Severino, C., & Lord, S.R. (2010). Effect on falls of providing single lens distance vision glasses to multifocal glasses wearers: VISIBLE Randomised Controlled Trial. *British Medical Journal*, 25, 340. c2265. doi: 10.1136/bmj.c2265.

Hawley-Hague, H., Horne, M., Campbell, M., Demack, S., Skelton, D. A., & Todd, C. (2013). Multiple levels of influence on older adults' attendance and adherence to community exercise classes. *The Gerontologist*, 54(4), 599–610. doi:10.1093/geront/gnt075

Horne, M., Skelton, D., Speed, S., & Todd, C. (2010). The influence of primary health care professionals in encouraging exercise and physical activity uptake among White and South Asian older adults: experiences of young older adults. *Patient Education and Counseling*, 78(1), 97–103. doi:10.1016/j.pec.2009.04.004

Laventure, R. M. E., Dinan, S. M., & Skelton, D. A. (2008). Someone like me: increasing participation in physical activity among seniors with senior peer health motivators. *Journal of Aging and Physical Activity*, (16), 76–7.

Fixsen, D., Scott. V., Blasé, K., Naoom, S., & Wagar, L. (2011). When evidence is not enough: the challenge of implementing fall prevention strategies. *Journal of Safety Research*, 42(6), 419-22. doi: 10.1016/j.jsr.2011.10.002

Sherrington, C., Whitney, J. C., Lord, S. R., Herbert, R. D., Cumming, R. G., & Close, J. C. T. (2008). Effective exercise for the prevention of falls: a systematic review and meta-analysis. *Journal of the American Geriatrics Society*, 56(12), 2234–2243. doi:10.1111/j.1532-5415.2008.02014.x

Yardley, L., Donovan-Hall, M., Francis, K., & Todd, C. (2007). Attitudes and beliefs that predict older people's intention to undertake strength and balance training. *The Journals of Gerontology. Series B, Psychological Sciences and Social Sciences*, 62(2), P119–125.

Yardley, L., Kirby, S., Ben-Shlomo, Y., Gilbert, R., Whitehead, S., & Todd, C. (2008). How likely are older people to take up different falls prevention activities? *Preventive Medicine*, 47(5), 554–558. doi:10.1016/j.ypmed.2008.09.001

Exercise

Many older people experience balance deficits and a reduction in muscular strength and power, due to ageing and to medical conditions and impairments that make movement less appealing. Yet, these are the two most important modifiable falls risk factors and can be influenced by a regular exercise training programme. Programmes must be tailored to the individual, be intensive enough to challenge balance and increase strength, progress in intensity and challenge over time, and last a sufficient duration, in order to be effective and to reduce the risk of falls.

What works?

Exercise is included in nearly all effective multiple interventions. To engage older people the emphasis should be primarily on strength and balance and healthy, active ageing and rather than falls prevention.

Programmes with multiple categories of exercise:

Multiple-component group and home-based exercise programmes (e.g. evidence based programmes such as Otago, FaME, LiFE), usually containing at least balance and strength training, have been proven to reduce falls. Other categories of exercise that can be included in multi-component exercise interventions include 3D training (constant repetitive unsupported movement through all three spatial planes) like Tai Chi and square stepping, general physical activity like walking groups, flexibility training or endurance training. The LiFE-programme comprises of balance and strength exercises embedded into daily activities of living and effectively reduces the rate of falls.

Exercise only interventions:

Multiple-component group exercise (eg. FaME) significantly reduces rate of falls and risk of falling, as does multiple-component home-based

exercise (eg. Otago, LiFE). For Tai Chi, the reduction in rate of falls bordered on statistical significance but Tai Chi does significantly reduce risk of falling.

The greatest relative effects of exercise on fall rates are seen in programmes that included a combination of a higher total dose of exercise (greater than 50 hours over the trial period) and challenging balance exercises (exercises conducted while standing in which people aimed to stand with their feet closer together or on one leg, minimize use of their hands to assist, and practice controlled movements of the center of mass) and did not include a walking program.

Multiple Interventions:

A study with “multifaceted” podiatry (customised orthoses, footwear review, falls prevention education), including foot and ankle exercises has been demonstrated to be as effective for preventing falls in older people with disabling foot pain.

Who can help older people with exercise?

Physiotherapists, sport scientists and specialist exercise instructors, who are appropriately trained in delivering falls prevention exercise programmes.

Assessment tools

Participants should be carefully assessed before intervention to ensure the correct type of programme is chosen and that the programme is tailored to their needs.

Appropriate assessment tools should be chosen to show progress, eg.

- Berg Balance Scale to assess balance
- Timed Up and Go to assess balance and mobility
- Short Physical Performance Battery to assess balance and strength
- Senior Fitness Test to assess balance, strength and endurance
- FES-I to assess fear of falling

What does not work?

There is no evidence for chair-based exercises in reducing falls. Brisk walking is not recommended for those at high risk of falls and can increase risk of falls for older people. Programmes that are only delivered for a short period of time may increase confidence without sufficiently improving strength and balance and reducing risk.

Summary

In order to be effective, exercise programmes must be challenging, progressive, at sufficient dosage and continued over time, they should:

- Focus on improving muscle strength/power of the lower limbs, ankles and feet
- Challenge balance in a standing position and/or gait (eg sideways, backwards walking)
- Exercise should be progressive and tailored to participants needs (help them to meet specific goals they have set, designed to consider health conditions).
- Be carried out 2-3 times a week
- At least 50 hours of strength and balance exercise should be carried out over a minimum of 3 months. Ideally exercise should be continued for maintenance of reduced risk.
- Be delivered by instructors specially trained in one of the following evidence based programmes (Regular contact and feedback from the instructor is helpful):

Evidence based programmes

Otago Exercise Programme

For the exercise booklet in a variety of languages please go to:

profound.eu.com/otago-exercise-program

For further information on the Otago Home Exercise Programme, visit www.cdc.gov/homeandrecreationalafety/pdf/cdc_falls_compendium_lowres.pdf

For information about Cascade Training in Otago across Europe, visit profound.eu.com/about/wp5-best-practice-exercise-regimen-net-work-development

Falls Management Exercise programme (FaME)

For the exercise booklet in a variety of languages please go to: profound.eu.com/strength-and-balance-home-exercise-booklet-for-older-people-english

For further information on the FaME exercise programme, visit www.cdc.gov/homeandrecreationalafety/pdf/cdc_falls_compendium_lowres.pdf

Or www.laterlifetraining.co.uk/fame-rationale-for-an-exercise-programme-to-prevent-falls

Lifestyle integrated Functional Exercise (LiFE)

profound.eu.com/life-lifestyle-integrated-functional-exercise-reducing-falls-and-improving-function

Square stepping

For more information about this programme including the evidence base please go to: square-step.org/en/home.html

Tai Chi

You can find out more about Tai Chi, Taijiquan and Qigong across Europe, visit taiji-europa.eu and www.tcfe.org

Links and Resources

Links related to exercise and falls

profound.eu.com

[www.ageuk.org.uk/Documents/EN-GB/For-professionals/Research/Falls Prevention Guide 2013.pdf](http://www.ageuk.org.uk/Documents/EN-GB/For-professionals/Research/Falls_Prevention_Guide_2013.pdf)

Other resources related to exercise and falls

Gait, balance and functional training

<http://profound.eu.com/video-clip-of-stronger-seniors-balance-exercise-program-english/>

Strength/resistance training

profound.eu.com/exercises-online-strengthening-video-english

Square Stepping Exercise

www.youtube.com/watch?v=lfCD7qB2l1k

Assessments

profound.eu.com/three-simple-assessment-tests-to-assess-the-patients-risk-for-falling

References

Gillespie, L. D., Robertson, M. C., Gillespie, W. J., Sherrington, C., Gates, S., Clemson, L. M., & Lamb, S. E. (2012). Interventions for preventing falls in older people living in the community. *The Cochrane Database of Systematic Reviews*, 9, CD007146. doi:10.1002/14651858.CD007146.pub3

Sherrington, C., Tiedemann, A., Fairhall, N., Close, J.C., & Lord, S.R. (2011). Exercise to prevent falls in older adults: an updated meta-analysis and best practice recommendations. *N S W Public Health Bull.* 22(3-4):78-83. doi: 10.1071/NB10056.

Spink, M.J., Menz, H.B., Fotoohabadi, M.R., Wee, E., Landorf, K.B., Hill, K.D., & Lord, S.R. (2011). Effectiveness of a multifaceted podiatry intervention to prevent falls in community dwelling older people with disabling foot pain: Randomised Controlled Trial. *British Medical Journal*, 16,342. d3411. doi: 10.1136/bmj.d3411.

Vision and falls

Impaired or low vision is an independent risk factor for falls in older people and unfamiliar glasses with a new vision correction or impaired vision affected by medication can increase the risk of older people falling. Many eye diseases such as cataract, age related macular degeneration, glaucoma and vascular eye disease are common in older people. Age related loss of contrast sensitivity and depth perception can cause balance problems.

When regular wearers of multifocal glasses were given single lens glasses falls were significantly reduced in the subgroup that regularly took part in outside activities. Conversely, there was a significant increase in out- side falls in intervention group participants who took part in little outside activity. New environments and a change in glasses or first prescription of glasses can increase the risk of falls during the first weeks because of altered and unfamiliar vision. This highlights the education role that optometrists and ophthalmologists can play.

Cataracts have been associated with increased risk of falls and fall related injury. First eye cataract surgery is a successful treatment that has been shown to reduce the risk of falling and fall related injuries.

Age related macular degeneration (AMD) is the most common form of age related vision loss in Europe. Currently there is no curative treatment for dry AMD, however compensating strategies have been tested. In people with very low vision and macular degeneration home visits by occupational therapists have been shown to reduce falls. A key element is the adaptation of the home environment and behavioural strategies for safe negotiation of the internal and external environment.

For patients suffering from glaucoma, diabetes and/or vascular eye disease or vision loss such as hemianopia no specific fall prevention programs have been tested so far. The treatment should be coordinated using best clinical practice and vision rehabilitation principles.

What works?

- Identifying new visual problems and ensuring spectacles are appropriate by testing visual acuity and glasses prescription every year
- Cataract surgery
- Occupational therapy
- Home safety and behavioural modifications for older people with very low vision such as AMD

Caution

- During the first days and weeks after a vision correction (e.g. new or changed prescription of eyeglasses or following cataract surgery) risk of falls can increase
- Using multifocal or bifocal lenses in new environments can increase the risk of falls
- There can be side effects of medication affecting visual acuity or adaptation

Who can help older people with their vision

- Ophthalmologists
- Optometrists
- Occupational therapist

Assessment tools

- Eye chart for testing visual acuity (e.g. Snellen chart)
- Melbourne Edge Test (MET) for testing contrast sensitivity

Links and Resources

profound.eu.com

References

Gillespie, L.D., Robertson, M.C., Gillespie, W.J., Sherrington, C., Gates, S., Clemson, L.M., & Lamb, S.E. (2012). Interventions for preventing falls in older people living in the community. *The Cochrane Database of Systematic Reviews*, 9, CD007146. doi:10.1002/14651858.CD007146.pub3

Lord, S.R., Smith, S., & Menant, C.J. (2010). Vision and Falls in Older People: Risk Factors and Intervention Strategies. *Clinics in Geriatric Medicine*, 26(4):569-81. doi: 10.1016/j.cger.2010.06.002.

Bone health

In older people, bone health is an important risk factor for fractures. Osteopenia and Osteoporosis, characterised by low bone mineral density (BMD), lead to bone fragility. Low vitamin D status is associated with a variety of negative skeletal consequences including osteomalacia, Reduced BMD, impaired calcium-absorption and secondary hyperparathyroidism. Falls risk should be taken into consideration when assessing whether or not to commence medication for osteoporosis. Bone health and fracture risk should be considered in a falls assessment.

What works?

Vitamin D (see factsheet vitamin D) and calcium supplementation is effective in reducing bone loss, falls and osteoporotic fractures:

- Vitamin D (1000-2000 IU/day; target levels: 50 nmol/L)
- Calcium (1000 to 1500 mg/day; if possible by nutrition intake, rather than tablets)

Nutrition/ lifestyle:

- Adequate calorie intake (Body Mass Index > 20 kg/m²)
- 1 g/kg body weight of protein per day
- Sufficient nutritional intake of Vitamin B12 (2,4 µg/day) and folic acid (400 µg/day)
- Cessation from smoking

Consideration of withdrawal of medication altering bone-health:

- Such as, glucocorticoids, glitazones, proton-pump inhibitors and antiepileptics in consultation with your doctor

Evidence-based anti-osteoporotic drugs (Bisphosphonates, denosumab, strontium ranelate, parathyroid hormone peptides):

Reduce the risk of vertebral fracture when given with calcium and vitamin D supplements.

Recommended for (see clinical guidelines in resources section or local Osteoporosis guidelines):

- Women with a prior fragility fracture, who should be considered for treatment without the need for further risk assessment
- Fragility vertebral fractures (single level 2 or 3 [moderate to severe], multiple level 1-3 [mild to severe])
- Fragility pertrochanteric fracture
- Fragility femoral neck fracture and T-score < -2,0 SD
- Therapy with glucocorticoids over > 3 month and T-score < -1,5 SD or fragility vertebral fractures
- Femoral neck T-score \leq to -2,5 SD (dependent on age and gender)

Identification and treatment of secondary causes of bone loss e.g. hypogonadism (low testosterone levels), reducing glucocorticoid medications, reducing alcohol consumption

Bone-health exercise programmes combined with fall prevention/exercise (see factsheet Exercise and Falls):

- Meta-analysis results suggest a relatively small statistically significant, but possibly important, effect of exercise on bone density compared with control groups. Exercise has the potential to be a safe and effective way to avert bone loss and reduce the risk of fracture
- Require a duration of 12 month and weight-bearing components to show effects on bone mineral density

- Weight-bearing components are successful in proven osteoporosis and / or after fractures
- Weight-bearing and resistance training with challenging balance exercises enhance bone and muscle health and improve functional ability
- A combination of weight-bearing impact exercise (jogging, stair-climbing, jumping activities) and progressive resistance training (PRT) is effective for maintaining bone mineral density (BMD) and preventing bone loss at clinically relevant sites such as the hip and spine

Who can help older people with impaired bone health?

- General physician, geriatrician, bone health specialist / endocrinologists/specialist osteoporosis nurse
- Consulting pharmacist
- Physiotherapists, sport scientists and exercise instructors, who are appropriately trained in delivering bone health and falls prevention exercise programs

Assessment tools

- European and/or national guidance for osteoporosis
- Bone mineral density (BMD) using dual-energy X-ray absorptiometry (DXA)
- X-ray (thoracic and lumbar spine) if vertebral fractures are suspected after clinical examination
- World Health Organization's Fracture Risk Assessment Tool (FRAX®)
- QFracture®, which uses falls risk in the algorithm
- Blood testing for differential diagnosis and specific treatment

Caution

Non-adherence to treatment with specific anti-osteoporosis drugs is a substantial problem. Medication can be inconvenient to take and unpleasant. Recommendation to monitor adherence.

Links and Resources

Links related to bone-health

www.iofbonehealth.org/europe-guidelines

The WHO fracture risk assessment tool:

www.shef.ac.uk/FRAX

QFracture risk tool:

www.qfracture.org

www.nice.org.uk/guidance/ta160

www.nice.org.uk/guidance/cg146

Also see guidelines on osteoporosis collected at

profound.eu.com/guidelines/

References

Edwards, M.H., Jameson, K., Denison, H., Harvey, N.C., Sayer, A.A., Dennison, E.M., & Cooper, C. (2013). Clinical risk factors, bone density and fall history in the prediction of incident fracture among men and women. *Bone*, 52(2), 541–547. doi:10.1016/j.bone.2012.11.006.

Gomez, F., Curcio, C. L., Suriyaarachchi, P., Demontiero, O., & Duque, G. (2013). Differing approaches to falls and fracture prevention between Australia Health Quality Ontario. (2008). Prevention of falls and fall-related injuries in community-dwelling seniors: an evidence-based analysis. *Ontario Health Technology Assessment Series*, 8(2), 1–78.

Kanis, J.A., McCloskey, E.V., Johansson, H., Cooper, C., Rizzoli, R., & Reginster, J-Y. (2013). European guidance for the diagnosis and management of osteoporosis in postmenopausal women. *Osteoporos Int*, 24(1), 23-57. DOI 10.1007/s00198-012-2074-y

Rizzoli, R., Bruyere, O., Cannata-Andia, J. B., Devogelaer, J.-P., Lyritis, G., Ringe, J. D., Vella, B., & Reginster, J-Y. (2009). Management of osteoporosis in the elderly. *Current Medical Research and Opinion*, 25(10), 2373–2387. doi:10.1185/03007990903169262

Howe, T.E., Shea, B., Dawson, L.J., Downie, F., Murray, A., Ross, C., Harbour, R.T., Caldwell, L.M. Creed, G. (2011). Exercise for preventing and treating osteoporosis in postmenopausal women. *The Cochrane Database of Systematic Reviews*, Issue 7. Art. No.: CD000333. doi: 10.1002/14651858.CD000333.pub2

Vitamin D

Low blood levels of vitamin D are associated with falls and some fall-related fractures. In many industrialised countries older people have modest to severe vitamin D deficiency. More than 70% of people in their 80s living in northern Europe suffer from vitamin D deficiency. This is caused by a combination of factors, such as nutritional deficiency, reduced renal function and skin atrophy. It is aggravated by a reduced sun exposure either through being housebound or institutionalised, seasonal lack of sun exposure due to weather (especially autumn/ winter) and lack of sunlight because of cultural factors (clothing which covers the skin from the sun). Severe vitamin D deficiency causes myopathy, loss of muscle strength and reduced bone health. Moderate deficiencies are linked to osteoporosis and impaired balance.

What works?

Taking vitamin D supplements may be effective in reducing falls in people with low blood levels of vitamin D. Maintaining adequate vitamin D levels, especially during winter, can be achieved through:

- Adequate sun exposure (face and arms), without sunscreen, of about 30 min in the middle of the day (depending on clothing, skin type, latitude and season). Longer periods should be avoided to limit the risk of skin cancer.
- Adequate nutritional intake from cod liver oil and fatty fish such as salmon, tuna or mackerel, beef liver, eggs, sardines and mushrooms
- When endogenous synthesis is missing (vitamin D from sun exposure), adequate vitamin D intake is estimated as 800 IU per day
- Supplementation with cholecalciferol for those at high risk of deficiency may require higher doses and should be discussed with the physicians.
- Supplementation can be taken as a weekly dose in drop form or a daily dose as a sweet. Both options are particularly suitable for older people in long-term care

Caution

- Overdosage of cholecalciferol is possible (but rare), leading to hypercalcaemia.
- Calcium levels must be monitored after four weeks to identify hypercalcaemia and possible hyperparathyroidism
- Large doses of cholecalciferol should not exceed 30,000 IU per week and should not be given for more than 10 weeks for a total of 300,000 IU.
- Check for contraindications
- Relevant reduction of dermal vitamin D synthesis through sun protection.

Who can help older people with their low vitamin D levels

- General physician
- Geriatrician
- Bone health specialist /endocrinologist
- Consulting pharmacist

Assessment tools

Blood levels of vitamin D (Target: Serum-25-Hydroxy-Vitamin D > 20 ng/ml (50 nmol))

References

American Geriatrics Society workgroup on Vitamin D supplementation of older adults. (2014). Recommendations Abstracted from the American Geriatrics Society Consensus Statement on Vitamin D for Prevention of Falls and Their Consequences. *Journal of American Geriatric Society*, 62 (1), 147–152. doi: 10.1111/jgs.12631

Avenell, A., Gillespie, W.J., Gillespie, L.D., & O'Connell, D. (2009). Vitamin D and vitamin analogues for preventing fractures associated with involutional and post-menopausal osteoporosis. *The Cochrane Database Systematic Review*, Issue 2. Art. No.: CD000227. doi.org/10.1002/14651858.CD000227.pub4

Bischoff- Ferrari, H.A., Dietrich, T., Orav, E.J., Hu, F.B., Zhang, Y., Karlson, E.W., & Dawson-Hughes, B. (2004). Higher 25-hydroxyvitamin D concentrations are associated with better lower-extremity function in both active and inactive persons aged > or =60 y. *The American Journal of Clinical Nutrition*. 80:752-758.

Gillespie, L.D., Robertson, M.C., Gillespie, W.J., Sherrington, C., Gates, S., Clemson, L.M., & Lamb, S.E. (2012). Interventions for preventing falls in older people living in the community. *The Cochrane Database of Systematic Review*, Issue 9, CD007146. doi:10.1002/14651858.CD007146.pub3

Osteoporosis Australia. (2013). Vitamin D position statement.

<http://www.osteoporosis.org.au/sites/default/files/files/Vit%20D%20Position%20Statement%2010%202013%20V2.pdf>

Home & environment

Numerous home and environmental factors, identified from reports on older fallers and structured observations, have been associated with falls in older people. Recent research considers environmental hazards as a modifiable external factor. The key factor is the relationship between the person's physical competence level and environmental stressors, e.g. an older person with some level of compromise of competence will fall at a lower level of perturbation than a younger or fitter older person. The level of risk of environmental hazards is influenced by lifestyle, risk taking, behaviour and exposure to environmental stressors, e.g. exposure to icy pavements, rooms with low lighting.

Some home safety assessment and modification interventions, alongside behavioural intervention, have been demonstrated to be effective in reducing both number of falls and number of people falling. Home safety assessment and modification appears to be effective for people at higher risk of falling (previous history of falls), particularly those with severe visual impairment. In a multifactorial intervention the home modification component should be implemented by occupational therapists and other trained specialist after a careful assessment of the personal environment.

Important components of environmental modifications (what works)

- Adjustments to ergonomic height of furniture (e.g. bed, chair, toilet seat)
- Adequate stability of furniture
- Barrier-free homes
- Handrails /holds
- Even and non-slip floorcoverings with optimised levels of friction
- Type of surface on which an older person falls, as this may reduce fall-related injuries (e.g. carpeted floors)

- Sufficient light and contrast in dark areas and at night, particularly on stairs
- Removal of trip hazards
- Urban planning considering the needs of older people (places to rest, even pathways/ pavements, ramps etc.)

What did not work?

- Home safety assessment and modification alone / as a single component intervention. (Effective interventions have been accompanied by education, training with transfers and provision of mobility aids).

Caution

- There is little high-level scientific evidence for modification of the built home environment as a method of reducing the risk of injury.
- Active people living with more environmental hazards are more likely to fall than frail people. Risk of falling is even increased in active people depending on their perception of risk and their outside participation. Frail people are more likely to fall because of their own limitations.
- Compliance issues are also caused by the stigmatising effects of home modifications (e.g. grab rails), and the perceived challenge to health and independence. Low financial or education status can also impact on older people's understanding of the need for home modification, limiting implementation (community and education programmes may help).

Who can help older people with home safety

- Occupational therapists (interventions appear to be more effective when carried out by an occupational therapist)
- Other trained specialist

Assessment tools

Home hazard checklists, such as Westmead Home Safety Assessment, HomeFast, EnableAge

Links related to home and environment

Westmead assessment tool

<https://sites.google.com/site/studentcapstone122/westmead-home-safety>

HomeFast assessment tool

www.bhps.org.uk/falls/documents/HomeFast.pdf

EnableAge assessment tool

www.enableage.arb.lu.se/pub.html

References

Lord, S. R., Sherrington, C., Menz, H. B., & Close, J. C. T. (2007). *Falls in Older People. Risk Factors for Prevention.* Cambridge: Cambridge University Press.

Gillespie, L. D., Robertson, M. C., Gillespie, W. J., Sherrington, C., Gates, S., Clemson, L. M., & Lamb, S. E. (2012). Interventions for preventing falls in older people living in the community. *The Cochrane Database of Systematic Reviews*, 9, CD007146. doi:10.1002/14651858.CD007146.pub3

Turner, S., Arthur, G., Lyons, R. A., Weightman, A. L., Mann, M. K., Jones, S. J., Lannon, S. (2011). Modification of the home environment for the reduction of injuries. *The Cochrane Database of Systematic Reviews*, (2), CD003600. doi:10.1002/14651858.CD003600.pub3

Cumming, R.G., Thomas, M., Szonyi, G., Salkeld, G., O'Neill, E., Westbury, C., Frampton, G. (1999). Home visits by an occupational therapist for assessment and modification of environmental hazards: a randomized trial of falls prevention. *Journal of the American Geriatrics Society*. 47(12):1397–1402.

Footwear & protective clothing

Footwear is very personal and can be culturally determined. Inappropriate footwear can increase the risk of slips, trips and falling and some certain shoes can decrease the base of support. An increased risk of falls has been linked to poor fitting shoes, slippers with a lack of heel support and high heeled shoes. Walking barefoot or with only stockings or socks indoors is also associated with an increased risk of falls.

Hip protectors can reduce the risk of fracture for older people living in nursing or residential care settings but has little effect for community dwelling older people due to low compliance.

What works?

- Footwear counselling- where older people are counselled on the specific identified hazardous features of their footwear and are provided with a handout on what constitutes a safe shoe.
- There is some evidence for the use of a non-slip device (Yaktrax® walker) on outdoor shoes in hazardous winter conditions.
- There is a reduction in rate of falls in people for people with disabling foot pain receiving “multifaceted podiatry” (customised orthoses, footwear review, foot and ankle exercises, fall prevention education, and “usual podiatry care”).
- Protective clothing such as hip protectors for older persons with high risk of fracture

Caution

Inappropriate footwear has been defined as a heel height >4.5 cm, or any two of the following:

- no fixation
- no heel counter (plastic or leather insert at the heel upper of the shoe)
- a heel counter that could be compressed greater than 45 degrees
- a fully worn or smooth sole
- or a shoe heel width narrower than the participant's heel by at least 20%.

Older people may not want to wear appropriate footwear or hip protectors because they do not fit with their personal identity. Having to wear sensible and practical footwear or hip protectors can identify them as 'a faller' and may cause issues with self-esteem and stigma.

Who can help older people with footwear

- Podiatry services
- Orthopedic specialist for severe deformities
- Trained nursing staff

References

Gillespie, L.D., Robertson, M.C., Gillespie, W.J., Sherrington, C., Gates, S., Clemson, L.M., & Lamb, S. E. (2012). Interventions for preventing falls in older people living in the community. *The Cochrane Database of Systematic Reviews*, 9, CD007146. doi:10.1002/14651858.CD007146.pub3

Lord, S. R., Sherrington, C., Menz, H. B., & Close, J. C. T. (2007). Falls in Older People. *Risk Factors for Prevention*. Cambridge: Cambridge University Press.

Spink, M., Menz, H.B., Fotoohabadi, M.R., Wee, E., Landorf, K.B., Hill, K.D., & Lord, S.R. (2011). Effectiveness of a multifaceted podiatry intervention to prevent falls in community-dwelling older people with disabling foot pain: a randomised controlled trial. *British Medical Journal*, 342:d3411. doi:10.1136/bmj.d3411.

Menz, H.B., & Sherrington, C. (2000). The Footwear Assessment Form: a reliable clinical tool to assess footwear characteristics of relevance to postural stability in older adults. *Clinical Rehabilitation*, 14 (6), 657-664. doi: 10.1191/0269215500cr375oa

Parker, M. J., Gillespie, W. J., & Gillespie, L. D. (2005). Hip protectors for preventing hip fractures in older people. *The Cochrane Database of Systematic Reviews*, (3), CD001255. doi:10.1002/14651858.CD001255.pub3

Santesso, N., Carrasco-Labra, A., & Brignardello-Petersen, R. (2014). Hip protectors for preventing hip fractures in older people. *The Cochrane Database of Systematic Reviews*, Issue 3. Art. No.: CD001255. doi: 10.1002/14651858.CD001255.pub5.

Falls detection and prevention technologies


There are a number of Information and Communication Technologies (ICTs) aimed at falls prevention, fall detection and alarms for use in case of a fall, that can alert professionals or carers if a fall occurs or if falls risk increases. There are also a range of ICT interventions which have been created or adapted to be pro-active in preventing falls, such as those which provide strength and balance training to older adults in the prevention of falls e.g. exergames, Wii-fit, Kinect.

What works?

- Personal Emergency Response Systems can help to reduce a long lie (lying on the floor for long periods of time can cause serious health complications) and allow help to be called quickly to the person who has fallen. However, there is evidence that people often do not press the alarms for fear of being a burden or nuisance. There is an emergence of automatic alarms that do not require the person to press them.
- Simple touchscreen interfaces and other easy to use technologies have been more readily accepted than those that are more complex and multi-faceted.
- Focusing on the possibility of maintaining independence is more likely to lead to successful uptake of PERS and ICT-devices.

Caution

- This is an emerging area and the evidence is currently not robust and is lacking completely in many areas, but this is changing rapidly.
- The main issues with use of ICT devices in the home are related to adoption and use of the systems; older adults need to understand the



value and potential of the technologies on offer and receive suitable training and support in using them.

- Evidence is weak around the use of virtual reality and gaming systems for the promotion of physical activity. Emerging evidence suggests that games should be designed specifically for and with older adults.
- Evidence is weak around the effective use of bed alarms in hospitals.

Who can help older people with ICT

- Occupational therapists/physiotherapists
- Social services and social care providers
- Sheltered and assisted housing providers
- Providers of telecare and telehealth equipment and services

References

Brownsell, S., & Hawley, M. S. (2004). Automatic fall detectors and the fear of falling. *Journal of Telemedicine and Telecare*, 10(5), 262–266.

FARSEEING Deliverable D5.2. “Validation strategy of the user interfaces, the fall risk assessment service & the exercise guidance service.” (n.d.). Retrieved from http://farseeingresearch.eu/wp-content/uploads/2014/02/D5-2_Final.pdf

Hawley-Hague, H., Boulton, E., Hall, A., Pfeiffer, K., & Todd, C. (2014). Older adults’ perceptions of technologies aimed at falls prevention, detection or monitoring: a systematic review. *International Journal of Medical Informatics*, 83(6), 416–426. doi:10.1016/j.ijmedinf.2014.03.002

Miller, K. J., Adair, B. S., Pearce, A. J., Said, C. M., Ozanne, E., & Morris, M. M. (2014). Effectiveness and feasibility of virtual reality and gaming system use at home by older adults for enabling physical activity to improve health-related domains: a systematic review. *Age and Ageing*, 43(2), 188–195. doi:10.1093/ageing/aft19

Schoene, D., Valenzuela, T., Lord, S.R., & de Bruin, E.D. (2014). The effect of interactive cognitive-motor training in reducing fall risk in older people: A systematic review. *BMC Geriatrics*, 14:107, doi:10.1186/1471-2318-14-107.

Shorr, R.I., Chandler, A.M., Mion, L.C., Waters, T.M., Minzhao Liu, M.S., Daniels, M.J., Kessler, L.A., & Miller, S.T.(2012). Effects of an intervention to increase bed alarm use to prevent falls in hospitalized patients: a cluster randomized trial. *Annals of Internal Medicine*. 157:692-699. doi: [10.7326/0003-4819-157-10-201211200-00005](https://doi.org/10.7326/0003-4819-157-10-201211200-00005)